

CSC 412/2506:
Probabilistic Learning and Reasoning
Week 12-1: Gaussian Processes

Murat A. Erdogdu

University of Toronto

Overview of the first hour

- We build on the kernel viewpoint of regression.
- We introduce Gaussian processes.
- This provides an additional component to kernel regression.
- We dispense with the parametric model and define a prior distribution over functions directly.
- There are multiple advantages (e.g. uncertainty quantification).

Recap: Linear Regression

- Given a training set of inputs and targets $\{(\mathbf{x}^{(i)}, y^{(i)})\}_{i=1}^N$
- Linear model:

$$y = \mathbf{w}^\top \boldsymbol{\psi}(\mathbf{x}) + \epsilon$$

where $\boldsymbol{\psi}(\mathbf{x})$ is the feature map.

- Vectorized, we have the design matrix \mathbf{X} in input space and

$$\boldsymbol{\Psi} = \begin{bmatrix} - & \boldsymbol{\psi}(\mathbf{x}^{(1)}) & - \\ - & \boldsymbol{\psi}(\mathbf{x}^{(2)}) & - \\ & \vdots & \\ - & \boldsymbol{\psi}(\mathbf{x}^{(N)}) & - \end{bmatrix} \in \mathbb{R}^{N \times M}$$

and predictions

$$\hat{\mathbf{y}} = \boldsymbol{\Psi} \mathbf{w}.$$

Recap: Bayesian Linear Regression

- We gave linear regression a probabilistic interpretation by assuming a Gaussian noise model:

$$y | \mathbf{x} \sim \mathcal{N}(\hat{y}(\mathbf{x}), \sigma^2), \quad \hat{y}(\mathbf{x}) = \mathbf{w}^\top \boldsymbol{\psi}(\mathbf{x})$$

- and a Gaussian prior

$$\mathbf{w} \sim \mathcal{N}\left(0, \frac{1}{\alpha} \mathbf{I}_M\right)$$

- Prior induces a probability distribution over

$$\hat{\mathbf{y}} = \boldsymbol{\Psi} \mathbf{w} \sim \mathcal{N}\left(0, \frac{1}{\alpha} \boldsymbol{\Psi} \boldsymbol{\Psi}^\top\right).$$

Distribution over prediction function

- In practice, we evaluate the prediction function $\hat{y}(\mathbf{x})$ at specific points, for example at the training data points $\mathbf{x}^{(i)}$ for $i = 1, \dots, N$.
- So we are interested in the joint distribution of the function values

$$\hat{y}(\mathbf{x}^{(1)}), \dots, \hat{y}(\mathbf{x}^{(N)})$$

which we denote by the vector $\hat{\mathbf{y}} = (\hat{y}(\mathbf{x}^{(1)}), \dots, \hat{y}(\mathbf{x}^{(N)}))$.

- We showed that

$$\hat{\mathbf{y}} \sim \mathcal{N}(0, \mathbf{K}) \quad \mathbf{K} = \frac{1}{\alpha} \mathbf{\Psi} \mathbf{\Psi}^\top$$

where \mathbf{K} is the (scaled) Gram matrix

$$K_{ij} = \frac{1}{\alpha} k(\mathbf{x}^{(i)}, \mathbf{x}^{(j)}) = \frac{1}{\alpha} \boldsymbol{\psi}(\mathbf{x}^{(i)})^\top \boldsymbol{\psi}(\mathbf{x}^{(j)})$$



Gaussian process

- **Definition:** A **Gaussian process** is a probability distribution over functions $\hat{y}(\mathbf{x})$ such that for any $N \geq 1$ and any set of N points $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(N)}$ in \mathbb{R}^D , the vector $(\hat{y}(\mathbf{x}^{(1)}), \dots, \hat{y}(\mathbf{x}^{(N)}))$ is jointly Gaussian.
- The joint distribution is specified completely by the second-order statistics, i.e. the mean and the covariance functions.
- In most applications, the mean function of $\hat{y}(\mathbf{x})$ can be set to zero and then the Gaussian process is completely specified by the covariance function

$$\mathbb{E}[\hat{y}(\mathbf{x})\hat{y}(\mathbf{x}')] = \frac{1}{\alpha}k(\mathbf{x}, \mathbf{x}')$$

Gaussian process

- We can directly define the kernel of a Gaussian process, not worrying about the feature map.



Samples from Gaussian processes for a Gaussian kernel (left) and an exponential kernel (right).

Gaussian processes for regression

- We have the linear model

$$y \mid \mathbf{x} \sim \mathcal{N}(\hat{y}(\mathbf{x}), \sigma^2) \quad \hat{y}(\mathbf{x}) = \mathbf{w}^\top \boldsymbol{\psi}(\mathbf{x})$$

- Given N independent observations, we have

$$\mathbf{y} \mid \hat{\mathbf{y}} \sim \mathcal{N}(\hat{\mathbf{y}}, \sigma^2 \mathbf{I}_N), \quad \hat{\mathbf{y}} \sim \mathcal{N}(0, \mathbf{K}).$$

- Therefore the marginal of \mathbf{y} is given by

$$\mathbf{y} \sim \mathcal{N}(0, \mathbf{C}) \quad \mathbf{C} = \mathbf{K} + \sigma^2 \mathbf{I}_N$$

where the corresponding kernel is

$$c(\mathbf{x}^{(i)}, \mathbf{x}^{(j)}) = \frac{1}{\alpha} k(\mathbf{x}^{(i)}, \mathbf{x}^{(j)}) + \sigma^2 \delta(\mathbf{x}^{(i)}, \mathbf{x}^{(j)})$$

$\delta(\mathbf{x}, \mathbf{x}') = 1$ if $\mathbf{x} = \mathbf{x}'$ and $\delta(\mathbf{x}, \mathbf{x}') = 0$ otherwise.

Gaussian processes for regression

- Denote now $\mathbf{y}_N = (y^{(1)}, y^{(2)}, \dots, y^{(N)})$.
- We have the marginal of \mathbf{y}_N given by

$$\mathbf{y}_N \sim \mathcal{N}(0, \mathbf{C}_N) \quad \mathbf{C}_N = \mathbf{K}_N + \sigma^2 \mathbf{I}_N.$$

- This reflects the two Gaussian sources of randomness.

Goal: We want to predict for a new output $y^{(N+1)}$.

- We need

$$p(y^{(N+1)} | \mathbf{y}_N)$$

- Note that $\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(N)}$ are treated as constants.

Gaussian processes for regression

- We have

$$\mathbf{y}_{N+1} \sim \mathcal{N}(0, \mathbf{C}_{N+1}) \quad \mathbf{C}_{N+1} = \mathbf{K}_{N+1} + \sigma^2 \mathbf{I}_{N+1}$$

where

$$\mathbf{C}_{N+1} = \begin{bmatrix} \mathbf{C}_N & \mathbf{k} \\ \mathbf{k}^\top & c \end{bmatrix}.$$

- ▶ Here, $c = \frac{1}{\alpha} k(\mathbf{x}^{(N+1)}, \mathbf{x}^{(N+1)}) + \sigma^2$
- ▶ \mathbf{k} is a vector with entries $k_i = \frac{1}{\alpha} k(\mathbf{x}^{(i)}, \mathbf{x}^{(N+1)})$
- Since the vector \mathbf{y}_{N+1} is Gaussian, we easily find $y^{(N+1)} | \mathbf{y}_N$.

Property of Multivariate Gaussian Distribution

Recall:

- If we have $\mathbf{x} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ with

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix} \quad \boldsymbol{\mu} = \begin{bmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{bmatrix} \quad \boldsymbol{\Sigma} = \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} \end{bmatrix}$$

- Then,

$$\mathbf{x}_2 \mid (\mathbf{x}_1 = \mathbf{a}) \sim \mathcal{N}(\mathbf{m}, \mathbf{C})$$

with

$$\mathbf{m} = \boldsymbol{\mu}_2 + \boldsymbol{\Sigma}_{21} \boldsymbol{\Sigma}_{11}^{-1} (\mathbf{a} - \boldsymbol{\mu}_1) \quad \mathbf{C} = \boldsymbol{\Sigma}_{22} - \boldsymbol{\Sigma}_{21} \boldsymbol{\Sigma}_{11}^{-1} \boldsymbol{\Sigma}_{12}.$$

Gaussian processes for regression

Recall:


$$\mathbf{y}_{N+1} \sim N(0, C_{N+1}), \quad C_{N+1} = \begin{bmatrix} C_N & \mathbf{k} \\ \mathbf{k}^\top & c \end{bmatrix}.$$

- Since \mathbf{y}_{N+1} is multivariate Gaussian, $y^{(N+1)} | \mathbf{y}_N$ is also Gaussian with mean and variance

$$\text{mean} = \mathbf{k}^\top C_N^{-1} \mathbf{y}_N \quad \text{variance} = c - \mathbf{k}^\top C_N^{-1} \mathbf{k}$$

- These are the key results that define Gaussian process regression.
- The vector \mathbf{k} is a function of the new test input $\mathbf{x}^{(N+1)}$.
- The predictive distribution is a Gaussian whose mean and variance both depend on $\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(N)}, \mathbf{x}^{(N+1)}$.


GPs for regression



- The green curve is the true sinusoidal function from which the data points, shown in blue, are obtained.
- The red line shows the **mean** of the Gaussian process predictive distribution.
- The shaded region corresponds to plus and minus two standard deviations.

GPs for classification

- Consider a classification problem with target variables $y \in \{0, 1\}$
- We define a Gaussian process over a function $a(\mathbf{x})$ and then transform the function using sigmoid $\hat{y}(\mathbf{x}) = \sigma(a(\mathbf{x}))$.
- We obtain a non-Gaussian stochastic process over functions $\hat{y}(\mathbf{x}) \in (0, 1)$.



Left: $a(\mathbf{x})$ Right: $\hat{y}(\mathbf{x})$

GPs for classification

- The probability distribution over target is then given by

$$p(y|a) = \sigma(a)^y(1 - \sigma(a))^{1-y}, \quad y \in \{0, 1\}.$$

- We need to compute

$$p(y^{(N+1)} | \mathbf{y}_N)$$

and notice that $a(\mathbf{x})$ is a Gaussian process but $\hat{y}(\mathbf{x})$ is not.

- We have $\mathbf{a}_{N+1} \sim \mathcal{N}(0, \mathbf{C}_{N+1})$, where

$$C_{N+1}(\mathbf{x}^{(i)}, \mathbf{x}^{(j)}) = \frac{1}{\alpha} k(\mathbf{x}^{(i)}, \mathbf{x}^{(j)}) + \nu \delta_{ij}.$$


- But \mathbf{a}_N is not observed, so we write

$$p(y^{(N+1)} | \mathbf{y}_N) = \int p(y^{(N+1)} | \mathbf{a}_{N+1}) p(\mathbf{a}_{N+1} | \mathbf{y}_N) d\mathbf{a}_{N+1}$$

- This is intractable. We need MCMC based methods, or numerical integration to approximate this integral.

GPs for classification: Illustration

- Illustration of GPs for classification:



- Left: optimal decision boundary from the true distribution in green, and the decision boundary from the Gaussian process classifier in black.
- Right: predicted posterior for the blue and red classes together with the Gaussian process decision boundary.

Learning the hyperparameters

- We didn't do any learning other than choosing a kernel!
- Rather than fixing the covariance function $\frac{1}{\alpha}k(\mathbf{x}, \mathbf{x}')$, we may prefer to use a parametric family of functions and then infer the parameter values from the data.
- Denoting the hyperparameters with θ , one can easily write down the likelihood of the Gaussian process model.

$$\log p(\mathbf{y} | \theta) = -\frac{1}{2} \log |\mathbf{C}_N| - \frac{1}{2} \mathbf{y}^\top \mathbf{C}_N^{-1} \mathbf{y} - \frac{N}{2} \log(2\pi)$$

- The next step is standard: gradient based optimization, grid search etc.

Summary of the first hour

- Gaussian processes are flexible tools that can be used in regression and classification tasks.
- One can simply choose a kernel and find the predictive density!
- They can be used together with modern tools, creating powerful learning methods.